

## A gated quantum dot strongly coupled to an optical microcavity

Systems

By Terra Quantum AG terraquantum.swiss

## A gated quantum dot strongly coupled to an optical microcavity

Daniel Najer, et al.

## ABSTRACT

The strong-coupling regime of cavity quantum electrodynamics (QED) represents the light-matter interaction at the fully quantum level. Adding a single photon shifts the resonance frequencies—a profound nonlinearity. Cavity QED is a test bed for quantum optics<sup>1,2,3</sup> and the basis of photon-photon and atom-atom entangling gates<sup>4,5</sup>. At microwave frequencies, cavity QED has had a transformative effect<sup>6</sup>, enabling qubit readout and qubit couplings in superconducting circuits. At optical frequencies, the gates are potentially much faster; the photons can propagate over long distances and can be easily detected. Following pioneering work on single atoms<sup>1,2,3,7</sup>, solid-state implementations using semiconductor quantum dots are emerging<sup>8,9,10,11,12,13,14,15</sup>. However, miniaturizing semiconductor cavities without introducing charge noise and scattering losses remains a challenge. Here we present a gated, ultralow-loss, frequency-tunable microcavity device. The gates allow both the quantum dot charge and its resonance frequency to be controlled electrically. Furthermore, cavity feeding<sup>10,11,13,14,15,16,17</sup>, the observation of the bare-cavity mode even at the quantum dot-cavity resonance, is eliminated. Even inside the microcavity, the quantum dot has a linewidth close to the radiative limit. In addition to a very pronounced avoided crossing in the spectral domain, we observe a clear coherent exchange of a single energy quantum between the 'atom' (the quantum dot) and the cavity in the time domain (vacuum Rabi oscillations), whereas decoherence arises mainly via the atom and photon loss channels. This coherence is exploited to probe the transitions between the singly and doubly excited photon-atom system using photon-statistics spectroscopy<sup>18</sup>. The work establishes a route to the development of semiconductorbased quantum photonics, such as single-photon sources and photon-photon gates.

The full article can be found here: https://www.nature.com/articles/s41586-019-1709-y